

Appln. No. 10/600,695
Response dated November 10, 2005
Reply to Non-Compliant Amendment dated October 27, 2005

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings of claims in the application:

Listing of Claims:

1. - 20. (Canceled)

21. (Currently amended) A pharmaceutical composition for preventing ~~or treating~~ a condition in a mammal characterized by undesired thrombosis comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of ~~claim 1~~ the formula (I):

A-Q-D-E-G-J-X

wherein:

A is selected from the group consisting of:

-C(=NR²)N(R²,R³); and

phenyl, which is substituted with 0-2 R¹ groups;

each R¹ is a member independently selected from the group consisting of:

halo, -CN, -C(=O)-N(R², R³), -NO₂, -SO₂N(R², R³), -SO₂R², -(CH₂)_mNR²R³, -(CH₂)_m-C(=NR³)-R², -(CH₂)_m-C(=NR²)-N(R²,R³), -(CH₂)_m-N(R²)-C(=NR²)-N(R²,R³), -(CH₂)_mNR²-C₃₋₆heterocyclics, C₁₋₄alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₈cycloalkyl, C₀₋₄alkylC₃₋₈cycloalkyl, -CF₃, -OR², and a 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C₁₋₄alkyl, -CN C₁₋₄alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₈cycloalkyl, C₀₋₄alkylC₃₋₈cycloalkyl and -NO₂;

each R² and R³ is a member independently selected from the group consisting of:

-H, -C₁₋₆alkyl, -C₁₋₆alkyloxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -S(=O)₂-OH, -CN, -CF₃ and -NO₂;

m is an integer of 0-2;

Q is a direct link;

D is phenyl, which is substituted with 0-2 R^{1a} groups;

each R^{1a} is a member independently selected from the group consisting of:

halo, -C₁₋₆alkyl, -C₁₋₆alkyloxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl, -S(=O)₂-OH, -CN, -NO₂, -(CH₂)_n-N(-R^{2a}, -R^{3a}), -S(=O)₂-N(-R^{2a}, -R^{3a}), -S(=O)₂-R^{2a}, -CF₃, -(CH₂)_n-OR^{2a}, -C(=O)-O-R^{2a}, -C(=O)-N(-R^{2a}, -R^{3a}), -C(=NH)-N(-R^{2a}, -R^{3a}), -C(=NMe)-N(-R^{2a}, -R^{3a}), 2-imidazolin-2-yl, 1-methyl-2-imidazolin-2-yl and a 5-6 membered aromatic heterocyclic ring containing 1-4 heteroatoms selected from N, O and S and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the aromatic heterocyclic ring and the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -CN, -CF₃ and -NO₂;

n is an integer of 0-2;

R^{2a} and R^{3a} are independently selected from the group consisting of:

-H, -C₁₋₆alkyl, -C₁₋₆alkyloxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -S(=O)₂-OH, -CN, -CF₃ and -NO₂;

E is -NH-C(=O)-;

G is a pyrazole ring substituted with 0-2 R^{1b} groups;

each R^{1b} is a member independently selected from the group consisting of:

halo, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl, -C₁₋₄alkyl-C(=O)-OH, -CN, -NO₂, -S(=O)₂-OH, -N(-R^{2b}, -R^{3b}), -C(=O)-N(-R^{2b}, -R^{3b}), -S(=O)₂-N(-R^{2b}, -R^{3b}), -S(=O)₂-R^{2b}, -CF₃, -O-R^{2b}, -O-CH₂-CH₂-O-R^{2b}, -O-CH₂-C(=O)-O-R^{2b}, -N(-R^{2b})-CH₂-CH₂-O-R^{2b}, -N(-CH₂-CH₂-O-R^{2b})₂, -N(-R^{2b})-C(=O)-R^{3b}, -N(-R^{2b})-S(=O)₂-R^{3b}, and a 5-6 membered heterocyclic ring containing 1-4 heteroatoms selected from N, O and S substituted with 0-4 R^{1b'} groups;

each R^{2b} and R^{3b} is a member independently selected from the group consisting of:

-H, -C₁₋₆alkyl, -C₁₋₆alkyloxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -S(=O)₂-O⁻, -CN, -CF₃ and -NO₂;

each R^{1b'} is a member independently selected from the group consisting of:

halo, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl, -C₁₋₄alkyl-C(=O)-OH, -CN, -NO₂, -S(=O)₂-OH, -N(-R^{2b'}, -R^{3b'}), -C(=O)-N(-R^{2b'}, -R^{3b'}), -S(=O)₂-N(-R^{2b'}, -R^{3b'}), -S(=O)₂-R^{2b'}, -CF₃, -O-R^{2b'}, -O-CH₂-CH₂-O-R^{2b'}, -O-CH₂-C(=O)-O-R^{2b'}, -N(-R^{2b'})-CH₂-CH₂-O-R^{2b'}, -N(-CH₂-CH₂-O-R^{2b'})₂, -N(-R^{2b'})-C(=O)-R^{3b'} and -N(-R^{2b'})-S(=O)₂-R^{3b'};

each R^{2b'} and R^{3b'} are independently selected from the group consisting of:

-H, -C₁₋₆alkyl, -C₁₋₆alkoxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -S(=O)₂-OH, -CN, -CF₃ and -NO₂;

J is a direct link;

X is a naphthyl, which is substituted with 0-3 R^{1c} groups;

each R^{1c} is a member independently selected from the group consisting of:

halo, -CF₃, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl, -C₁₋₄alkyl-C(=O)-OH, -CF₃, -CN, -NO₂, -(CH₂)_z-N(-R^{2c}, -R^{3c}), -C(=O)-N(-R^{2c}, -R^{3c}), -C(=NH)-N(-R^{2c}, -R^{3c}), -C(=NMe)-N(-R^{2c}, -R^{3c}), -S(=O)₂-N(-R^{2c}, -R^{3c}), -S(=O)₂-R^{2c}, -S(=O)₂-OH, -CF₃, -O-R^{2c}, -O-(CH₂)_z-O-R^{2c}, -O-(CH₂)_z-C(=O)-O-R^{2c}, -N(-R^{2c}), -O-(CH₂)_z-O-R^{2c}, -N[(CH₂)_z-O-R^{2c}]₂, -(CH₂)_z-N(-R^{2c})-C(=O)-R^{3c}, -(CH₂)_z-N(-R^{2c})-S(=O)₂-R^{3c}, and a 5-6 membered heterocyclic ring containing 1-4 heteroatoms selected from N, O and S;

z is an integer of 0-4;

each R^{2c} and R^{3c} is a member independently selected from the group consisting of:

-H, -C₁₋₆alkyl, -C₁₋₆alkyloxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -S(=O)₂-OH, -CN, -CF₃ and -NO₂;

and all pharmaceutically acceptable diastereomers, enantiomers or mixtures thereof, salts, hydrates or solvates thereof.

22. (Currently amended) A method for preventing ~~or treating~~ a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of ~~claim 1~~ the formula (I):

A-Q-D-E-G-J-X

wherein:

A is selected from the group consisting of:

-C(=NR²)N(R²,R³); and

phenyl, which is substituted with 0-2 R¹ groups;

each R¹ is a member independently selected from the group consisting of:

halo, -CN, -C(=O)-N(R², R³), -NO₂, -SO₂N(R², R³), -SO₂R², -(CH₂)_mNR²R³, -(CH₂)_m-C(=NR³)-R², -(CH₂)_m-C(=NR²)-N(R²,R³), -(CH₂)_m-N(R²)-C(=NR²)-N(R²,R³), -(CH₂)_mNR²-C₃₋₆heterocyclics, C₁₋₄alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₈cycloalkyl, C₀₋₄

alkylC₃₋₈cycloalkyl, -CF₃, -OR², and a 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C₁-C₄-alkyl, -CN C₁₋₄alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₈cycloalkyl, C₀₋₄alkylC₃₋₈cycloalkyl and -NO₂;

each R² and R³ is a member independently selected from the group consisting of:

-H, -C₁₋₆alkyl, -C₁₋₆alkyloxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -S(=O)₂-OH, -CN, -CF₃ and -NO₂;

m is an integer of 0-2;

Q is a direct link;

D is phenyl, which is substituted with 0-2 R^{1a} groups;

each R^{1a} is a member independently selected from the group consisting of:

halo, -C₁₋₆alkyl, -C₁₋₆alkyloxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl, -S(=O)₂-OH, -CN, -NO₂, -(CH₂)_n-N(-R^{2a}, -R^{3a}), -S(=O)₂-N(-R^{2a}, -R^{3a}), -S(=O)₂-R^{2a}, -CF₃, -(CH₂)_n-OR^{2a}, -C(=O)-O-R^{2a}, -C(=O)-N(-R^{2a}, -R^{3a}), -C(=NH)-N(-R^{2a}, -R^{3a}), -C(=NMe)-N(-R^{2a}, -R^{3a}), 2-imidazolin-2-yl, 1-methyl-2-imidazolin-2-yl and a 5-6 membered aromatic heterocyclic ring containing 1-4 heteroatoms selected from N, O and S and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the aromatic heterocyclic ring and

the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -CN, -CF₃ and -NO₂;

n is an integer of 0-2;

R^{2a} and R^{3a} are independently selected from the group consisting of:

-H, -C₁₋₆alkyl, -C₁₋₆alkyloxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -S(=O)₂-OH, -CN, -CF₃ and -NO₂;

E is -NH-C(=O)-;

G is a pyrazole ring substituted with 0-2 R^{1b} groups;

each R^{1b} is a member independently selected from the group consisting of:

halo, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl, -C₁₋₄alkyl-C(=O)-OH, -CN, -NO₂, -S(=O)₂-OH, -N(-R^{2b}, -R^{3b}), -C(=O)-N(-R^{2b}, -R^{3b}), -S(=O)₂-N(-R^{2b}, -R^{3b}), -S(=O)₂-R^{2b}, -CF₃, -O-R^{2b}, -O-CH₂-CH₂-O-R^{2b}, -O-CH₂-C(=O)-O-R^{2b}, -N(-R^{2b})-CH₂-CH₂-O-R^{2b}, -N(-CH₂-CH₂-O-R^{2b})₂, -N(-R^{2b})-C(=O)-R^{3b}, -N(-R^{2b})-S(=O)₂-R^{3b}, and a 5-6 membered heterocyclic ring containing 1-4 heteroatoms selected from N, O and S substituted with 0-4 R^{1b'} groups;

each R^{2b} and R^{3b} is a member independently selected from the group consisting of:

-H, -C₁₋₆alkyl, -C₁₋₆alkyloxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -S(=O)₂-O⁻, -CN, -CF₃ and -NO₂;

each $R^{1b'}$ is a member independently selected from the group consisting of:

halo, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl, -C₁₋₄alkyl-C(=O)-OH, -CN, -NO₂, -S(=O)₂-OH, -N($R^{2b'}$, $R^{3b'}$), -C(=O)-N($R^{2b'}$, $R^{3b'}$), -S(=O)₂-N($R^{2b'}$, $R^{3b'}$), -S(=O)₂- $R^{2b'}$, -CF₃, -O- $R^{2b'}$, -O-CH₂-CH₂-O- $R^{2b'}$, -O-CH₂-C(=O)-O- $R^{2b'}$, -N($R^{2b'}$)-CH₂-CH₂-O- $R^{2b'}$, -N(-CH₂-CH₂-O- $R^{2b'}$)₂, -N($R^{2b'}$)-C(=O)- $R^{3b'}$ and -N($R^{2b'}$)-S(=O)₂- $R^{3b'}$;

each $R^{2b'}$ and $R^{3b'}$ are independently selected from the group consisting of:

-H, -C₁₋₆alkyl, -C₁₋₆alkoxy, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₆alkylC₃₋₈cycloalkyl and -C₀₋₆alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, -C₁₋₄alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -C₃₋₈cycloalkyl, -C₀₋₄alkylC₃₋₈cycloalkyl, -S(=O)₂-OH, -CN, -CF₃ and -NO₂;

J is a direct link;

X is a naphthyl, which is substituted with 0-3 R^{1c} groups;

each R^{1c} is a member independently selected from the group consisting of:

halo, $-CF_3$, $-C_{1-6}$ alkyl, $-C_{2-6}$ alkenyl, $-C_{2-6}$ alkynyl, $-C_{3-8}$ cycloalkyl, $-C_{0-6}$ alkyl C_{3-8} cycloalkyl, $-C_{1-4}$ alkyl- $C(=O)$ -OH, $-CF_3$, $-CN$, $-NO_2$, $-(CH_2)_z-N(-R^{2c}, -R^{3c})$, $-C(=O)-N(-R^{2c}, -R^{3c})$, $-C(=NH)-N(-R^{2c}, -R^{3c})$, $-C(=NMe)-N(-R^{2c}, -R^{3c})$, $-S(=O)_2-N(-R^{2c}, -R^{3c})$, $-S(=O)_2-R^{2c}$, $-S(=O)_2-OH$, $-CF_3$, $-O-R^{2c}$, $-O-(CH_2)_z-O-R^{2c}$, $-O-(CH_2)_z-C(=O)-O-R^{2c}$, $-N(-R^{2c})$, $-O-(CH_2)_z-O-R^{2c}$, $-N[(CH_2)_z-O-R^{2c}]_2$, $-(CH_2)_z-N(-R^{2c})-C(=O)-R^{3c}$, $-(CH_2)_z-N(-R^{2c})-S(=O)_2-R^{3c}$, and a 5-6 membered heterocyclic ring containing 1-4 heteroatoms selected from N, O and S;

z is an integer of 0-4;

each R^{2c} and R^{3c} is a member independently selected from the group consisting of:

$-H$, $-C_{1-6}$ alkyl, $-C_{1-6}$ alkyloxy, $-C_{2-6}$ alkenyl, $-C_{2-6}$ alkynyl, $-C_{3-8}$ cycloalkyl, $-C_{0-6}$ alkyl C_{3-8} cycloalkyl and $-C_{0-6}$ alkyl-(carbocyclic aryl), wherein from 0-4 hydrogen atoms on the ring atoms of the carbocyclic aryl moiety may be independently replaced with a member selected from the group consisting of halo, $-C_{1-4}$ alkyl, $-C_{2-6}$ alkenyl, $-C_{2-6}$ alkynyl, $-C_{3-8}$ cycloalkyl, $-C_{0-4}$ alkyl C_{3-8} cycloalkyl, $-S(=O)_2-OH$, $-CN$, $-CF_3$ and $-NO_2$;

and all pharmaceutically acceptable diastereomers, enantiomers or mixtures thereof, salts, hydrates or solvates thereof.

23. (Currently amended) The method of claim 22 6, wherein the condition is selected from the group consisting of:

acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke,

thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation, and thrombotic complications associated with the fitting of prosthetic devices.

24. (Canceled)

25. (Currently amended) A pharmaceutical composition of claim 21 for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of claim 2

wherein:

A is phenyl, which is substituted with 0-2 R¹ groups;

each R¹ is a member independently selected from the group consisting of:

halo, C₁₋₄alkyl, -CN, -C(=O)-N(R², R³), -NO₂, -SO₂N(R², R³), -SO₂R², -(CH₂)_mNR²R³, -(CH₂)_m-C(=NR³)-R², -(CH₂)_m-C(=NR²)-N(R², R³), -(CH₂)_m-N(R²)-C(=NR²)-N(R², R³), -(CH₂)_mNR²-C₃₋₆heterocyclics, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₈cycloalkyl, C₀₋₄alkylC₃₋₈cycloalkyl, -CF₃, -OR², and a 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

each R² and R³ is a member independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₀₋₄alkyl-(carbocyclic aryl);

Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

m is an integer of 0-2;

Q is a direct link;

D is phenyl, which is substituted with 0-2 R^{1a} groups;

each R^{1a} is a member independently selected from the group consisting of:

halo, -C₁₄alkyl, -CN, -NO₂, -(CH₂)_n-N(-R^{2a}, -R^{3a}), -S(=O)₂-N(-R^{2a}, -R^{3a}), -S(=O)₂-R^{2a}, -CF₃, -(CH₂)_n-OR^{2a}, -C(=O)-O-R^{2a}, -C(=O)-N(-R^{2a}, -R^{3a}), and a 5-6 membered aromatic heterocyclic ring containing 1-4 heteroatoms selected from N, O and S;

n is an integer of 0-2;

R^{2a} and R^{3a} are independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₁₋₄alkyl-(carbocyclic aryl);

E is -NH-C(=O)-;

G is a pyrazole ring substituted with 0-2 R^{1b} groups;

each R^{1b} is a member independently selected from the group consisting of:

halo, -C₁₋₄alkyl, -CN, -NO₂, -N(-R^{2b}, -R^{3b}), -C(=O)-N(-R^{2b}, -R^{3b}), -S(=O)₂-N(-R^{2b}, -R^{3b}), -S(=O)₂-R^{2b}, -CF₃, -O-R^{2b}, -O-CH₂-CH₂-O-R^{2b}, -O-CH₂-C(=O)-O-R^{2b}, -N(-R^{2b})-CH₂-CH₂-O-R^{2b}, -N(-CH₂-CH₂-O-R^{2b})₂, -N(-R^{2b})-C(=O)-R^{3b}, -N(-R^{2b})-S(=O)₂-R^{3b}, and a 5-6 membered heterocyclic ring containing 1-4 heteroatoms selected from N, O and S;

Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

each R^{2b} and R^{3b} is a member independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₁₋₄alkyl-(carbocyclic aryl);

J is a direct link;

X is a naphthyl, which is substituted with 0-3 R^{1c} groups;

each R^{1c} is a member independently selected from the group consisting of:

halo, -C₁₋₄alkyl, -CN, -NO₂, -(CH₂)_z-N(-R^{2c}, -R^{3c}), -C(=O)-N(-R^{2c}, -R^{3c}),
-C(=NH)-N(-R^{2c}, -R^{3c}), -C(=NMe)-N(-R^{2c}, -R^{3c}), -S(=O)₂-N(-R^{2c}, -R^{3c}), -S(=O)₂-R^{2c},
-S(=O)₂-OH, -CF₃, -O-R^{2c}, -O-(CH₂)_z-O-R^{2c}, -O-(CH₂)_z-C(=O)-O-R^{2c}, -N(-R^{2c}),
-O-(CH₂)_z-O-R^{2c}, -N[(CH₂)_z-O-R^{2c}]₂, -(CH₂)_z-N(-R^{2c})-C(=O)-R^{3c},
-(CH₂)_z-N(-R^{2c})-S(=O)₂-R^{3c}, and a 5-6 membered heterocyclic ring containing 1-4
heteroatoms selected from N, O and S;

z is an integer of 0-4;

each R^{2c} and R^{3c} is a member independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₁₋₄alkyl-(carbocyclic aryl);

or all pharmaceutically acceptable diastereomers, enantiomers or mixtures thereof, salts, hydrates or solvates thereof.

26. (Currently amended) ~~The A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 2~~ 22

wherein:

Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

A is selected from the group consisting of:

phenyl, which is substituted with 0-2 R¹ groups;

each R¹ is a member independently selected from the group consisting of:

halo, C₁₋₄alkyl, -CN, -C(=O)-N(R², R³), -NO₂, -SO₂N(R², R³), -SO₂R², -(CH₂)_mNR²R³, -(CH₂)_m-C(=NR³)-R², -(CH₂)_m-C(=NR²)-N(R², R³), -(CH₂)_m-N(R²)-C(=NR²)-N(R², R³), -(CH₂)_mNR²-C₃₋₆heterocyclics, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₈cycloalkyl, C₀₋₄alkylC₃₋₈cycloalkyl, -CF₃, -OR², and a 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

each R² and R³ is a member independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₀₋₄alkyl-(carbocyclic aryl);

m is an integer of 0-2;

Q is a direct link;

D is phenyl, which is substituted with 0-2 R^{1a} groups;

each R^{1a} is a member independently selected from the group consisting of:

halo, -C₁₋₄alkyl, -CN, -NO₂, -(CH₂)_n-N(-R^{2a}, -R^{3a}), -S(=O)₂-N(-R^{2a}, -R^{3a}), -S(=O)₂-R^{2a}, -CF₃, -(CH₂)_n-OR^{2a}, -C(=O)-O-R^{2a}, -C(=O)-N(-R^{2a}, -R^{3a}), and a 5-6 membered aromatic heterocyclic ring containing 1-4 heteroatoms selected from N, O and S;

n is an integer of 0-2;

R^{2a} and R^{3a} are independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₁₋₄alkyl-(carbocyclic aryl);

Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

E is -NH-C(=O)-;

G is a pyrazole ring substituted with 0-2 R^{1b} groups;

each R^{1b} is a member independently selected from the group consisting of:

halo, -C₁₋₄alkyl, -CN, -NO₂, -N(-R^{2b}, -R^{3b}), -C(=O)-N(-R^{2b}, -R^{3b}), -S(=O)₂-N(-R^{2b}, -R^{3b}), -S(=O)₂-R^{2b}, -CF₃, -O-R^{2b}, -O-CH₂-CH₂-O-R^{2b}, -O-CH₂-C(=O)-O-R^{2b}, -N(-R^{2b})-CH₂-CH₂-O-R^{2b}, -N(-CH₂-CH₂-O-R^{2b})₂, -N(-R^{2b})-C(=O)-R^{3b}, -N(-R^{2b})-S(=O)₂-R^{3b}, and a 5-6 membered heterocyclic ring containing 1-4 heteroatoms selected from N, O and S;

each R^{2b} and R^{3b} is a member independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₁₋₄alkyl-(carbocyclic aryl);

J is a direct link;

X is a naphthyl, which is substituted with 0-3 R^{1c} groups;

each R^{1c} is a member independently selected from the group consisting of:

halo, -C₁₋₄alkyl, -CN, -NO₂, -(CH₂)_z-N(-R^{2c}, -R^{3c}), -C(=O)-N(-R^{2c}, -R^{3c}), -C(=NH)-N(-R^{2c}, -R^{3c}), -C(=NMe)-N(-R^{2c}, -R^{3c}), -S(=O)₂-N(-R^{2c}, -R^{3c}), -S(=O)₂-R^{2c}, -S(=O)₂-OH, -CF₃, -O-R^{2c}, -O-(CH₂)_z-O-R^{2c}, -O-(CH₂)_z-C(=O)-O-R^{2c}, -N(-R^{2c}), -O-(CH₂)_z-O-R^{2c}, -N[-(CH₂)_z-O-R^{2c}]₂, -(CH₂)_z-N(-R^{2c})-C(=O)-R^{3c}, -(CH₂)_z-N(-R^{2c})-S(=O)₂-R^{3c}, and a 5-6 membered heterocyclic ring containing 1-4 heteroatoms selected from N, O and S;

z is an integer of 0-4;

Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

each R^{2c} and R^{3c} is a member independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₁₋₄alkyl-(carbocyclic aryl);

or all pharmaceutically acceptable diastereomers, enantiomers or mixtures thereof, salts, hydrates or solvates thereof.

27. (Currently amended) The method of claim ~~26~~ 10, wherein the condition is selected from the group consisting of:

acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation, and thrombotic complications associated with the fitting of prosthetic devices.

28. (Canceled)

29. (Currently amended) A pharmaceutical composition of claim 21 ~~for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of claim 3~~


wherein:

Appln. No. 10/600,695

Response dated November 10, 2005


Reply to Non-Compliant Amendment dated October 27, 2005

A is selected from the group consisting of:




Q is a direct link;

D is selected from the group consisting of:



E is $-\text{NH}-\text{C}(=\text{O})-$;

G has the following formula:



each R^{1b} is a member independently selected from the group consisting of:

$-\text{H}$, $-\text{Me}$, $-\text{CF}_3$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{SO}_2\text{Me}$, $-\text{CN}$, $-\text{CONH}_2$, $-\text{CONMe}_2$, $-\text{NH}_2$, $-\text{NO}_2$, $-\text{NHCOMe}$, $-\text{NHSO}_2\text{Me}$, $-\text{CH}_2\text{NH}_2$ and $-\text{CO}_2\text{H}$;


Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

J is a direct link;


X is selected from the group consisting of:





Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005







or all pharmaceutically acceptable diastereomers, enantiomers or mixtures thereof, salts, hydrates or solvates thereof

30. (Currently amended) The A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 3 22


wherein:

Appln. No. 10/600,695

Response dated November 10, 2005


Reply to Non-Compliant Amendment dated October 27, 2005

A is selected from the group consisting of:




Q is a direct link;

D is selected from the group consisting of:



E is -NH-C(=O)-;

G has the following formula:



each R^{1b} is a member independently selected from the group consisting of:

-H, -Me, -CF₃, -F, -Cl, -Br, -SO₂Me, -CN, -CONH₂, -CONMe₂, -NH₂, -NO₂, -NHCOMe, -NHSO₂Me, -CH₂NH₂ and -CO₂H;


Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

J is a direct link;


X is selected from the group consisting of:




Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005






Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005



or all pharmaceutically acceptable diastereomers, enantiomers or mixtures thereof, salts, hydrates or solvates thereof.

31. (Original) The method of claim 30, wherein the condition is selected from the group consisting of:

acute coronary syndrome, myocardial infarction, unstable angina, refractory angina,

occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation, and thrombotic complications associated with the fitting of prosthetic devices.

32. (Canceled)

33. (Currently amended) A pharmaceutical composition of claim 21 for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of claim 4

wherein:

A is phenyl, which is substituted with 0-2 R¹ groups;

each R¹ is a member independently selected from the group consisting of:

halo, C₁₋₄alkyl, -CN, -C(=O)-N(R², R³), -NO₂, -SO₂N(R², R³), -SO₂R², -(CH₂)_mNR²R³, -(CH₂)_m-C(=NR³)-R², -(CH₂)_m-C(=NR²)-N(R², R³), -(CH₂)_m-N(R²)-C(=NR²)-N(R², R³), -(CH₂)_mNR²-C₃₋₆heterocyclics, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₈cycloalkyl, C₀₋₄alkylC₃₋₈cycloalkyl, -CF₃, -OR², and a 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

each R^2 and R^3 is a member independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₀₋₄alkyl-(carbocyclic aryl);

m is an integer of 0-2;

Q is a direct link;

D is phenyl, which is substituted with 0-2 R^{1a} groups;

each R^{1a} is a member independently selected from the group consisting of:

halo, -C₁₋₄alkyl, -CN, -NO₂, -(CH₂)_n-N(-R^{2a}, -R^{3a}), -S(=O)₂-N(-R^{2a}, -R^{3a}), -S(=O)₂-R^{2a},
-CF₃, -(CH₂)_n-OR^{2a}, -C(=O)-O-R^{2a}, -C(=O)-N(-R^{2a}, -R^{3a}), and a 5-6 membered
aromatic heterocyclic ring containing 1-4 heteroatoms selected from N, O and S;

n is an integer of 0-2;

R^{2a} and R^{3a} are independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₁₋₄alkyl-(carbocyclic aryl);

E is -NH-C(=O)-;

G is a pyrazole ring substituted with 0-2 R^{1b} groups;

each R^{1b} is a member independently selected from the group consisting of:

halo, -C₁₋₄alkyl, -CN, -NO₂, -N(-R^{2b}, -R^{3b}), -C(=O)-N(-R^{2b}, -R^{3b}), -S(=O)₂-N(-R^{2b}, -
R^{3b}), -S(=O)₂-R^{2b}, -CF₃, -O-R^{2b}, -O-CH₂-CH₂-O-R^{2b}, -O-CH₂-C(=O)-O-R^{2b},
-N(-R^{2b})-CH₂-CH₂-O-R^{2b}, -N(-CH₂-CH₂-O-R^{2b})₂, -N(-R^{2b})-C(=O)-R^{3b},
-N(-R^{2b})-S(=O)₂-R^{3b}, and a 5-6 membered heterocyclic ring containing 1-4
heteroatoms selected from N, O and S;

Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

each R^{2b} and R^{3b} is a member independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₁₋₄alkyl-(carbocyclic aryl);

J is a direct link;

X is a naphthyl, which is substituted with 0-3 R^{1c} groups;

each R^{1c} is a member independently selected from the group consisting of:

halo, -C₁₋₄alkyl, -CN, -NO₂, -(CH₂)_z-N(-R^{2c}, -R^{3c}), -C(=O)-N(-R^{2c}, -R^{3c}),
-C(=NH)-N(-R^{2c}, -R^{3c}), -C(=NMe)-N(-R^{2c}, -R^{3c}), -S(=O)₂-N(-R^{2c}, -R^{3c}), -S(=O)₂-R^{2c},
-S(=O)₂-OH, -CF₃, -O-R^{2c}, -O(-CH₂)_z-O-R^{2c}, -O(-CH₂)_z-C(=O)-O-R^{2c}, -N(-R^{2c}),
-O(-CH₂)_z-O-R^{2c}, -N[(-CH₂)_z-O-R^{2c}]₂, -(CH₂)_z-N(-R^{2c})-C(=O)-R^{3c},
-(CH₂)_z-N(-R^{2c})-S(=O)₂-R^{3c}, and a 5-6 membered heterocyclic ring containing 1-4
heteroatoms selected from N, O and S;

z is an integer of 0-4;

each R^{2c} and R^{3c} is a member independently selected from the group consisting of:

-H, -C₁₋₄alkyl and -C₁₋₄alkyl-(carbocyclic aryl);

or all pharmaceutically acceptable diastereomers, enantiomers or mixtures thereof, salts, hydrates or solvates thereof.

34. (Currently amended) The A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 4 22

wherein:

Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

A is phenyl, which is substituted with 0-2 R¹ groups;

each R¹ is a member independently selected from the group consisting of:

halo, -CN, -SO₂N(R², R³), -SO₂R² and -CH₂NR²R³;

each R² and R³ is a member independently selected from the group consisting of:

-H and -C₁₋₄alkyl;

Q is a direct link;

D is phenyl, which is substituted with 0-2 R^{1a} groups;

each R^{1a} is a member independently selected from the group consisting of:

-H and halo;

E is -NH-C(=O)-;

G is a pyrazole ring substituted with 0-2 R^{1b} groups;

each R^{1b} is a member independently selected from the group consisting of:

- Me, -Et, -CF₃, -C(=O)-NH₂, -NH₂, -NH-(C=O)-Me, -NH-S(=O)₂-Me, -SMe, -S(=O)-

Me and halo;

J is a direct link;

X is a naphthyl, which is substituted with 0-3 R^{1c} groups;

each R^{1c} is a member independently selected from the group consisting of:

**halo, OH, -OMe, -NH₂, -CN, -NO₂, -CH₂OH, -C₁₋₅alkyl, -C(=O)-N(-R^{2c}, -R^{3c}),
-C(=NH)-N(-R^{2c}, -R^{3c}), -S(=O)₂-N(-R^{2c}, -R^{3c}), -S(=O)₂-R^{2c}, -S(=O)₂-OH, -CF₃, 2-
imidazolin-2-yl and 1-methyl-2-imidazolin-2-yl;**

each R^{2c} and R^{3c} is a member independently selected from the group consisting of:

-H, -OH, -NH₂ and -C₁₋₄alkyl;

or all pharmaceutically acceptable diastereomers, enantiomers or mixtures thereof, salts, hydrates or solvates thereof.


35. (Original) The method of claim 34, wherein the condition is selected from the group consisting of:

acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation, and thrombotic complications associated with the fitting of prosthetic devices.

36. (Canceled)

37. (New) A pharmaceutical composition of claim 21

wherein the compound has the following formula:



R^1 is selected from the group consisting of:

$-S(=O)_2-NH_2$, $-S(=O)_2-Me$, $-CH_2NH_2$, and $-CH_2NMe_2$;

R^{1a} is selected from the group consisting of:

$-H$, $-F$, $-Cl$ and $-Br$;

R^{1c1} is independently selected from the group consisting of:

$-H$, $-F$, $-Cl$, $-Br$, $-NH_2$, $-OH$, $-SO_2Me$, $-SO_2Et$, $-SO_2NH_2$, $-NO_2$, $-CN$, $-CONH_2$ and $-CH_2OH$;

R^{1c2} is independently selected from the group consisting of:

$-H$, $-F$, $-Cl$ and $-Br$;

R^{1c3} is independently selected from the group consisting of:

$-H$, $-F$, $-Cl$ and $-Br$;

Appln. No. 10/600,695

Response dated November 10, 2005


Reply to Non-Compliant Amendment dated October 27, 2005

R^{1b} is selected from the group consisting of:

-H, -CH₃ and -CF₃.

38. (New) The method of claim 22

wherein the compound has the following formula:



R¹ is selected from the group consisting of:

-S(=O)₂-NH₂, -S(=O)₂-Me, -CH₂NH₂, and -CH₂NMe₂;

R^{1a} is selected from the group consisting of:

-H, -F, -Cl and -Br;

R^{1c1} is independently selected from the group consisting of:

-H, -F, -Cl, -Br, -NH₂, -OH, -SO₂Me, -SO₂Et, -SO₂NH₂, -NO₂, -CN, -CONH₂ and -CH₂OH;

R^{1c2} is independently selected from the group consisting of:

-H, -F, -Cl and -Br;

Appln. No. 10/600,695

Response dated November 10, 2005

Reply to Non-Compliant Amendment dated October 27, 2005

R^{1c3} is independently selected from the group consisting of:

-H, -F, -Cl and -Br;

R^{1b} is selected from the group consisting of:

-H, -CH₃ and -CF₃.

39. (New) The method of claim 38, wherein the condition is selected from the group consisting of:

acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation, and thrombotic complications associated with the fitting of prosthetic devices.